

Geology

Linking orography, climate, and exhumation across the central Andes

Jason B. Barnes, Todd A. Ehlers, Nadja Insel, Nadine McQuarrie and Christopher J. Poulsen

Geology published online 18 September 2012;
doi: 10.1130/G33229.1

Email alerting services click www.gsapubs.org/cgi/alerts to receive free e-mail alerts when new articles cite this article

Subscribe click www.gsapubs.org/subscriptions/ to subscribe to *Geology*

Permission request click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Notes

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by GeoRef from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

Linking orography, climate, and exhumation across the central Andes

Jason B. Barnes¹, Todd A. Ehlers², Nadja Insel³, Nadine McQuarrie⁴, and Christopher J. Poulsen⁵

¹Department of Geological Sciences, University of North Carolina, Chapel Hill, North Carolina 27599, USA

²Department of Geosciences, University of Tübingen, Tübingen 72076, Germany

³Department of the Geophysical Sciences, University of Chicago, Chicago, Illinois 60637, USA

⁴Department of Geology and Planetary Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA

⁵Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, Michigan 48109-1005, USA

ABSTRACT

Quantifying interactions between uplift, climate, deformation, and exhumation remains difficult, in many cases due to a paucity of data relevant to all processes. We synthesize new and existing data to understand the orogen-scale orographic changes across the central Andes, Bolivia. We use a regional climate model and geo-thermochronologic data to identify the correlations between changes in precipitation due to surface uplift and spatiotemporal patterns of deformation and erosional exhumation. Mean orographic rainfall patterns do not reach near present-day gradients and values until the topography grows to >75% modern elevations. New fission-track data near the orocline apex indicate that rapid exhumation moved eastward, beginning in the Eastern Cordillera ca. 50–15 Ma, the Interandean zone ca. 18–6 Ma, and in the Subandes ca. 7–3 Ma. Throughout Bolivia, exhumation is consistent with deformation until ca. 15–11 Ma, after which the pattern corresponds better with the increased rainfall toward modern values. These linked observations suggest that ca. 15–11 Ma, regional elevations reached threshold values (>75% modern) necessary to generate near present-day, enhanced rainfall gradients. These gradients have resulted in variable exhumation implied by the structural level of rocks exposed across the thrust belt and confirmed by fission tracks in apatite. The main insight is that the climate-induced Middle Miocene–recent exhumation varies over scales of a few hundred kilometers across Bolivia and implies that high mean rainfall (>3 m/yr) and long time scales (~10 m.y.) may be necessary for climate to induce orographically driven exhumation patterns recorded by fission tracks.

INTRODUCTION

Interactions between tectonics, topography, climate, and erosion evolve as mountains grow. Numerical models suggest coupling of tectonics and climate (Willett, 1999) and dramatic climate shifts triggered by increased surface elevations (e.g., Prell and Kutzbach, 1992). However, the significance of these insights is limited by the tenuous connection between the models and data available to critically evaluate them (Whipple, 2009). This failing exists, in part, because data sets that address orogen evolution often

represent disparate scales. For example, time spans covered by geologic data can be long (one to tens of millions of years), but modern climate patterns may be short-lived (millennia), even though they are often assumed invariant over geologic time scales (Montgomery et al., 2001; Reiners et al., 2003).

Models show that Andean surface uplift strongly affects regional climate (Lenters and Cook, 1995). Orographic enhancement of precipitation patterns could result in variable orogen erosion recorded by exhumation patterns quan-

tified over million-year time scales using thermochronology. We focus on the central Andes because correlations in topography and climate suggest that erosion has influenced thrust belt development (Horton, 1999) and shortening and exhumation records advocate that tectonics and climate became connected during the Miocene (e.g., Strecker et al., 2007). We hypothesize that this connection developed as elevations reached heights sufficient to induce intense orographic precipitation and erosion gradients across the Bolivian orocline (Masek et al., 1994; Norton and Schlunegger, 2011). We test this idea with new mean rainfall estimates from climate simulations that are forced by rising Andean elevations and link them to the spatiotemporal patterns of exhumation constrained by geothermochronologic data.

GEOLOGIC SETTING

Eocene–present (ca. 45–0 Ma) central Andes mountain building has produced a retroarc plateau and thrust belt (Fig. 1). The thrust belt has migrated eastward through time and varies in width and shortening along strike (McQuarrie et al., 2008a). The plateau and thrust belt is divided into the Altiplano basin (AL) and the higher relief regions of the Eastern Cordillera (EC), Interandean zone (IA), and Subandes (SA). These zones step downward from the EC to the SA in mean elevation and upward in structural depth, exposing Paleozoic to Tertiary rocks (Kley, 1999). In Bolivia, the thrust belt




Figure 1. Central Andes topography, tectonics, and climate. Transects: N—north, C—center, S—south. A: Thrust belt region bounding faults and zones: AL—Altiplano, EC—Eastern Cordillera, IA—Interandean zone, SA—Subandes, FL—foreland. B: Tropical Rainfall Measuring Mission (TRMM) estimated rainfall (1998–2006; Bookhagen and Strecker, 2008). Insets are 50 km swath mean elevation (km; black line, gray is range; 90 m digital elevation model) and rainfall (m/yr; blue line \pm 20%) profiles.

straddles a sharp change in precipitation, from subtropical dry conditions south of ~18°S to subequatorial wet conditions to the north. Orography alters the rainfall distribution, producing small mean rainfall gradients (~2 to <1 m/yr) in the south compared to the center and north (~4 to <1 m/yr; Fig. 1B).

METHODS: CLIMATE MODEL AND THERMOCHRONOLOGY

REMO_{iso} is a limited-domain general circulation model (~55 km resolution) that replicates modern South American climate patterns well (Sturm et al., 2007). Topography is the most important factor forcing Andean flank climate gradients on geologic time scales (Jeffery et al., 2011). We extracted mean rainfall values from simulations with 50%, 75%, and 100% mean modern Andean elevations to examine the first-order response of rainfall patterns to surface uplift along three thrust belt transects (Fig. 2A) (Insel et al., 2012; see the GSA Data Repository¹).

Low-temperature thermochronology can resolve the ~50–350 °C cooling history of rocks to quantify regional patterns of orogen erosion (e.g., Gillis et al., 2006). We combine 17 new fission-track samples (see the Data Repository) that traverse the thrust belt apex with other transects of published data (~125 samples; Fig. 1). We interpret the data with standard methods of (1) thermal modeling to constrain cooling

histories and identify the onset of most recent, rapid exhumation and (2) estimating depths to closure temperatures using geothermal data to constrain the exhumation magnitude (cf. Barnes et al., 2006, 2008). For brevity, we summarize the constraints from good-fit thermal models associated with the robust apatite fission-track data (samples with >10 measured grains, >20 measured tracks; see the Data Repository) combined with published ⁴⁰Ar/³⁹Ar, fission-track, and (U-Th)/He data (Fig. 2B).

RESULTS AND INTERPRETATIONS

Climate simulations predict the response of rainfall to Andean surface uplift (Fig. 2A). At 50% modern elevations, mean rainfall rates decrease from ≤2 m/yr with increasing elevation across the northern and central transects and have no gradient across the southern transect. At 75% modern elevations, rates increase to 2.8 m/yr, but remain far below the observed peak values (Figs. 2A and 1B); only above 75% modern elevations are near modern rainfall maxima and orographic gradients established. Specifically, rainfall maxima (~3.8 m/yr or more) become focused in the northern EC and in the eastern IA to SA along the central transect. The control case (100% modern elevations) simulates the across-strike rainfall patterns well, but broadens the distribution of orographic precipitation and overpredicts the rainfall in places (e.g., the AL; Fig. 2A), common biases in most

climate models (Codron and Sadourny, 2002). Regardless, the precipitation sensitivity to uplift and the >75% modern elevation threshold inducing peak rainfall is consistent with many models of various resolutions and elevation gain increments, and is related to enhanced latent heat transport, convergence, and convection (Insel et al., 2009; Poulsen et al., 2010).

The Cenozoic history of central Andean growth shows that deformation and initial rapid exhumation are synchronous and that this initial exhumation front tracks the eastward propagation of the thrust belt from the EC since the Eocene (ca. 45 Ma; Fig. 2B) (Barnes and Ehlers, 2009). In the south, exhumation began in the central EC ca. 36–32 Ma, moved into the IA ca. 22–19 Ma, and into the SA ca. 20–8 Ma (Barnes et al., 2008). Results from further south at ~21°S are similar (Fig. 1A; Scheuber et al., 2006). In the center, exhumation began in the EC ca. 50–45 Ma, in the IA ca. 18–6 Ma, and in the SA ca. 7–3 Ma. In the north, exhumation began ca. 50–30 Ma in the EC, ca. 25 Ma or earlier in the IA, and ca. 20–8 in the SA (McQuarrie et al., 2008b). Collectively, deformation and initial exhumation migrated eastward across the EC ca. 50–30 Ma, the IA ca. 25–18 Ma, and the SA ca. 20–3 Ma.

The eastward propagation of linked deformation and exhumation breaks down along the northern transect, marking a vital change in the exhumation history ca. 15–11 Ma (Fig. 2B).




Figure 2. Climate and exhumation along transects in Figure 1. A: Regional circulation model REMO_{iso} rainfall estimates for 50%, 75%, and 100% modern (mod) elevations (elev) compared to 100% model elevations. Insets are ~150 km swath profiles for 100% mod elevation run for comparison to Figure 1B profiles. B: Cordilleran exhumation patterns (Fig. 1A data). Gray is rapid exhumation (exhum.) since ca. 15–11 Ma that correlates with climate in north. Inset is spatial extent overlying Tropical Rainfall Measuring Mission (TRMM) rainfall (Fig. 1B). Stars are best age constraints (age-elevation profiles) for both exhumation phases in the northern EC (Gillis et al., 2006).

¹GSA Data Repository item 2012328, climate model and fission-track data, methods, and results, is available online at www.geosociety.org/pubs/ft2012.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

Here, EC exhumation began at 40 Ma, best constrained by age-elevation profiles of biotite $^{40}\text{Ar}/^{39}\text{Ar}$ and zircon fission-track data (Benjamin et al., 1987; Gillis et al., 2006). A second rapid exhumation phase began ca. 15–11 Ma, spanning the entire EC to SA (Barnes et al., 2006). The best constraint (an apatite fission-track age-elevation profile) for this phase is 11 Ma, which postdates EC upper crustal deformation by ~14 m.y. (Gillis et al., 2006).

Thermochronology data also estimate exhumation magnitudes across the orogen (Fig. 2B). Precambrian–Mesozoic (ca. 600–150 Ma) zircon fission-track and (U–Th)/He ages limit Andean exhumation to <10–8 km in the southern and central EC–IA regions and to <7 km in the northern SA. In the south, Cenozoic apatite fission-track data suggest ~4 km of exhumation from the EC and IA and ~3 km from the SA. Results from ~21°S suggest ~6–2 km of exhumation across the EC and IA (Ege et al., 2007). In the center, we estimate ~4.4 km of exhumation from the EC and IA, and ~4 km from the SA. The north is different where Eocene–Oligocene $^{40}\text{Ar}/^{39}\text{Ar}$ ages imply ~10 km of total exhumation in the EC compared to ~4 km in the IA and SA (Gillis et al., 2006). These results show that exhumation has barely decreased eastward across the orogen, with the exception of the northern EC.

The SA apatite fission-track data record a south to north gradient in exhumation magnitude from low to high to intermediate, coincident with modern rainfall. This is evident from Paleozoic–Mesozoic cooling ages exhumed from partial annealing zone temperatures at ≤3.5 km in the south, reset Miocene–Pliocene cooling ages exhumed from below closure temperature at ≥4 km in the center, and mostly early Cenozoic mixed and/or partially reset cooling ages exhumed from partial annealing zone temperatures at ~4 km depth in the north (this study; Barnes et al., 2008).

In summary, climate simulations indicate that near modern rainfall gradients become established after >75% modern elevations are attained. The exhumation patterns emphasize a Middle Miocene (ca. 15–11 Ma) shift to distributed exhumation in the northern transect, unassociated with EC upper crustal deformation, yet ~4 km in magnitude since ca. 11 Ma (Gillis et al., 2006). The patterns also feature Middle Miocene–Pliocene (ca. 13–3 Ma) exhumation in the eastern IA–SA in the central transect and imply that SA exhumation magnitude varies along strike. These collective exhumation patterns coincide better with the modern rainfall patterns than they do with the eastward propagation of Miocene–recent deformation (e.g., gray; Fig. 2B).

Erosion is the main exhumation mechanism in active thrust belts, often facilitated by deformation creating topographic relief. How-

ever, exhumation can continue after deformation ceases and can vary in response to climate change (e.g., Fig. 2A). We propose that near modern rainfall gradients have persisted enough to affect the regional exhumation patterns, best exemplified by the northern transect with distributed exhumation since ca. 15–11 Ma and a high magnitude in the EC. Apatite fission-track data record the erosion of several kilometers of rock (sensitive to ~110 °C; geothermal gradients are ~35–20 °C/km; see the Data Repository). We propose that near-modern rainfall gradients were established at >75% modern elevations ca. 15–11 Ma and have mainly driven the Middle Miocene to present exhumation patterns north of 18°S.

DISCUSSION

Existing geologic and isotopic records in Bolivia suggest changes in climate and topography consistent with increasing rainfall and altitudes ca. 15–11 Ma (Fig. 3). SA stratigraphy records a tenfold increase in sedimentation rate with Yecua Formation deposition at 12.4 Ma (Uba et al., 2009). A decrease in $\delta^{18}\text{O}$ of pedogenic carbonate from these strata starting ca. 12 Ma is interpreted to indicate low $\delta^{18}\text{O}$ rainfall due to enhanced rainout caused by growing topography (Mulch et al., 2010). By ca. 9–6 Ma, increased climate seasonality and a sediment source change via headward catchment growth caused a further increase in

sedimentation and an increase in multiple soil carbonate isotope ratios (Mulch et al., 2010; Uba et al., 2009). The northern EC exhumation ca. 11 Ma is recorded by ongoing Cangalli Formation deposition ca. 10 Ma in an adjacent wedge-top basin, followed by its incision beginning ca. 7 Ma (Mosolf et al., 2010).

Critical taper theory for thrust belt growth offers a mechanical basis for erosion restricting orogen width (e.g., Suppe, 1981). The SA are a good test case because deformation began ca. 16 Ma or later, coincident with or after the establishment of the near modern rainfall gradients (Fig. 3) (Espurt et al., 2011). North to south variations in SA shortening (41%, 55%, 32%), width (96, 56, 139 km), and modern rainfall are consistent with erosion limiting thrust belt propagation along the northern and central transects relative to the south (Figs. 1 and 3) (McQuarrie, 2002; McQuarrie et al., 2008a). This suggests a tectonics–climate correlation at a scale of a few hundred kilometers. However, the shortening estimates may overlap within uncertainty and inherited tectonostratigraphic differences may also affect propagation (Judge and Allmendinger, 2011).

The northern EC and eastern IA to SA in the central transect are sites of high (~4 km) and young (Miocene–Pliocene) exhumation (Fig. 2B; Gillis et al., 2006). This is also implied by the (1) peak rainfall today (~4 m/yr; Fig. 1B), (2) deepest stratigraphic exposures,




Figure 3. Geologic data consistent with enhanced rainfall starting 15–11 Ma (dashed lines); sed—sedimentation, S—shortening. Transects: N—north, C—center, S—south; AL—Altiplano, EC—Eastern Cordillera, IA—Interandean zone, SA—Subandes. See text for discussion. Central Andes deformation history is integrated over all transects plus data from Peru and Argentina (Espurt et al., 2011). Inset is latitudinal variations in SA shortening with 5% error (McQuarrie, 2002; McQuarrie et al., 2008a).

i.e., Cambrian in the central SA only at ~17°S (McQuarrie, 2002) and lowermost Ordovician in the northern EC at ~16.5°S (McQuarrie et al., 2008b), and (3) high channel concavities signifying climatic forcing in the northern EC (Schlunegger et al., 2011). These results broadly suggest that high (~3 m/yr or more) mean orographic rainfall and ~10 m.y. time scales (~15–11 m.y. here) may be required to link fission-track–recorded exhumation and climate patterns in other active orogens.

ACKNOWLEDGMENTS

We thank Sohrab Tawackoli and Jaime Tito for logistical support and the National Science Foundation for financial support (grants EAR-0409289, EAR-0738822, and EAR-0907817 to Ehlers and Poulsen). We thank F. Schlunegger, J.-D. Champagnac, and an anonymous reviewer for their constructive reviews.

REFERENCES CITED

- Barnes, J.B., and Ehlers, T.A., 2009, End member models for Andean Plateau uplift: *Earth-Science Reviews*, v. 97, p. 105–132, doi:10.1016/j.earscirev.2009.08.003.
- Barnes, J.B., Ehlers, T.A., McQuarrie, N., O'Sullivan, P.B., and Pelletier, J.D., 2006, Variations in Eocene to recent erosion across the central Andean fold-thrust belt, northern Bolivia: Implications for plateau evolution: *Earth and Planetary Science Letters*, v. 248, p. 118–133, doi:10.1016/j.epsl.2006.05.018.
- Barnes, J.B., Ehlers, T.A., McQuarrie, N., O'Sullivan, P.B., and Tawackoli, S., 2008, Thermochronometer record of central Andean Plateau growth, Bolivia (19.5°S): *Tectonics*, v. 27, TC3003, doi:10.1029/2007TC002174.
- Benjamin, M.T., Johnson, N.M., and Naeser, C.W., 1987, Recent rapid uplift in the Bolivian Andes; evidence from fission-track dating: *Geology*, v. 15, p. 680–683, doi:10.1130/0091-7613(1987)15<680:RRUITB>2.0.CO;2.
- Bookhagen, B., and Strecker, M.R., 2008, Orographic barriers, high-resolution TRMM rainfall, and relief variations along the eastern Andes: *Geophysical Research Letters*, v. 35, L06403, doi:10.1029/2007GL032011.
- Codron, F., and Sadourny, R., 2002, Saturation limiters for water vapour advection schemes: Impact on orographic precipitation: *Tellus*, v. 54A, p. 338–349, doi:10.1034/j.1600-0870.2002.01265.x.
- Ege, H., Sobel, E.R., Scheuber, E., and Jacobshagen, V., 2007, Exhumation history of the southern Altiplano plateau (southern Bolivia) constrained by apatite fission-track thermochronology: *Tectonics*, v. 26, TC1004, doi:10.1029/2005TC001869.
- Espurt, N., Barbarand, J., Roddaz, M., Brusset, S., Baby, P., Saillard, M., and Hermoza, W., 2011, A scenario for late Neogene Andean shortening transfer in the Camisea Subandean zone (Peru, 12°S): Implications for growth of the northern Andean Plateau: *Geological Society of America Bulletin*, v. 123, p. 2050–2068, doi:10.1130/B30165.1.
- Gillis, R.J., Horton, B.K., and Grove, M., 2006, Thermochronology, geochronology, and upper crustal structure of the Cordillera Real: Implications for Cenozoic exhumation of the central Andean plateau: *Tectonics*, v. 25, TC6007, doi:10.1029/2005TC001887.
- Horton, B.K., 1999, Erosional control on the geometry and kinematics of thrust belt development in the central Andes: *Tectonics*, v. 18, p. 1292–1304, doi:10.1029/1999TC900051.
- Insel, N., Poulsen, C., and Ehlers, T., 2009, Influence of the Andes Mountains on South American moisture transport, convection, and precipitation: *Climate Dynamics*, v. 35, p. 1477–1492, doi:10.1007/s00382-009-0637-1.
- Insel, N., Poulsen, C.J., Ehlers, T.A., and Sturm, C., 2012, Response of meteoric $\delta^{18}\text{O}$ to surface uplift—Implications for Cenozoic Andean Plateau growth: *Earth and Planetary Science Letters*, v. 317–318, p. 262–272, doi:10.1016/j.epsl.2011.11.039.
- Jeffery, M.L., Poulsen, C.J., and Ehlers, T.A., 2011, Impacts of Cenozoic global cooling, surface uplift, and an inland seaway on South American paleoclimate and precipitation $\delta^{18}\text{O}$: *Geological Society of America Bulletin*, doi:10.1130/B30480.30481.
- Judge, P.A., and Allmendinger, R.W., 2011, Assessing uncertainties in balanced cross sections: *Journal of Structural Geology*, v. 33, p. 458–467, doi:10.1016/j.jsg.2011.01.006.
- Kley, J., 1999, Geologic and geometric constraints on a kinematic model of the Bolivian Orocline: *Journal of South American Earth Sciences*, v. 12, p. 221–235, doi:10.1016/S0895-9811(99)00015-2.
- Lenters, J.D., and Cook, K.H., 1995, Simulation and diagnosis of the regional summertime precipitation climatology of South America: *Journal of Climate*, v. 8, p. 2988–3005, doi:10.1175/1520-0442(1995)008<2988:SADOTR>2.0.CO;2.
- Masek, J.G., Isacks, B.L., Gubbels, T.L., and Fielding, E.J., 1994, Erosion and tectonics at the margins of continental plateaus: *Journal of Geophysical Research*, v. 99, p. 13,941–13,956, doi:10.1029/94JB00461.
- McQuarrie, N., 2002, The kinematic history of the central Andean fold-thrust belt, Bolivia: Implications for building a high plateau: *Geological Society of America Bulletin*, v. 114, p. 950–963, doi:10.1130/0016-7606(2002)114<0950:TKHOTC>2.0.CO;2.
- McQuarrie, N., Ehlers, T.A., Barnes, J.B., and Meade, B.J., 2008a, Temporal variation in climate and tectonic coupling in the central Andes: *Geology*, v. 36, p. 999–1002, doi:10.1130/G25124A.1.
- McQuarrie, N., Barnes, J.B., and Ehlers, T.A., 2008b, Geometric, kinematic, and erosional history of the central Andean Plateau, Bolivia (15–17°S): *Tectonics*, v. 27, TC3007, doi:10.1029/2006TC002054.
- Montgomery, D.R., Balco, G., and Willett, S.D., 2001, Climate, tectonics, and the morphology of the Andes: *Geology*, v. 29, p. 579–582, doi:10.1130/0091-7613(2001)029<0579:CTATMO>2.0.CO;2.
- Mosolf, J.G., Horton, B.K., Heizler, M.T., and Matos, R., 2010, Unroofing the core of the central Andean fold-thrust belt during focused late Miocene exhumation: Evidence from the Tipuani-Mapiri wedge-top basin, Bolivia: *Basin Research*, v. 23, p. 346–360, doi:10.1111/j.1365-2117.2010.00491.x.
- Mulch, A., Uba, C.E., Strecker, M.R., Schoenberg, R., and Chamberlain, C.P., 2010, Late Miocene climate variability and surface elevation in the central Andes: *Earth and Planetary Science Letters*, v. 290, p. 173–182, doi:10.1016/j.epsl.2009.12.019.
- Norton, K., and Schlunegger, F., 2011, Migrating deformation in the Central Andes from enhanced orographic rainfall: *Nature Communications*, v. 2, no. 584, doi:10.1038/ncomms1590.
- Poulsen, C.J., Ehlers, T.A., and Insel, N., 2010, Onset of convective rainfall during gradual late Miocene rise of the Central Andes: *Science*, v. 328, p. 490–493, doi:10.1126/science.1185078.
- Prell, W.L., and Kutzbach, J.E., 1992, Sensitivity of the Indian monsoon to forcing parameters and implications for its evolution: *Nature*, v. 360, p. 647–652, doi:10.1038/360647a0.
- Reiners, P.W., Ehlers, T.A., Mitchell, S.G., and Montgomery, D.R., 2003, Coupled spatial variations in precipitation and long-term erosion rates across the Washington Cascades: *Nature*, v. 426, p. 645–647, doi:10.1038/nature02111.
- Safran, E.B., Blythe, A., and Thomas, D., 2006, Spatially variable exhumation rates in orogenic belts: An Andean example: *Journal of Geology*, v. 114, p. 665–681, doi:10.1086/507613.
- Scheuber, E., Mertmann, D., Ege, H., Silva-Gonzalez, P., Heubeck, C., Reutter, K.-J., and Jacobshagen, V., 2006, Exhumation and basin development related to formation of the central Andean plateau, 21°S, *in* Oncken, O., et al., eds., *The Andes: Active subduction orogeny*: New York, Springer-Verlag, p. 285–301.
- Schlunegger, F., Norton, K.P., and Zeilinger, G., 2011, Climatic forcing on channel profiles in the Eastern Cordillera of the Coroico region, Bolivia: *Journal of Geology*, v. 119, p. 97–107, doi:10.1086/657407.
- Strecker, M.R., Alonso, R.N., Bookhagen, B., Carrapa, B., Hilley, G.E., Sobel, E.R., and Trauth, M.H., 2007, Tectonics and climate of the southern Central Andes: *Annual Review of Earth and Planetary Sciences*, v. 35, p. 747–787, doi:10.1146/annurev.earth.35.031306.140158.
- Sturm, C., Hoffmann, G., and Langmann, B., 2007, Simulation of the stable water isotopes in precipitation over South America: Comparing regional to global circulation models: *Journal of Climate*, v. 20, p. 3730–3750, doi:10.1175/JCLI4194.1.
- Suppe, J., 1981, Mechanics of mountain building and metamorphism in Taiwan: *Geological Society of China Memoir*, v. 4, p. 67–89.
- Uba, C.E., Kley, J., Strecker, M.R., and Schmitt, A., 2009, Unsteady evolution of the Bolivian Subandean thrust belt: The role of enhanced erosion and clastic wedge progradation: *Earth and Planetary Science Letters*, v. 281, p. 134–146, doi:10.1016/j.epsl.2009.02.010.
- Whipple, K.X., 2009, The influence of climate on the tectonic evolution of mountain belts: *Nature Geoscience*, v. 2, p. 97–104, doi:10.1038/ngeo413.
- Willett, S.D., 1999, Orogeny and orography; the effects of erosion on the structure of mountain belts: *Journal of Geophysical Research*, v. 104, p. 28,957–28,981, doi:10.1029/1999JB900248.

Manuscript received 23 January 2012

Revised manuscript received 25 May 2012

Manuscript accepted 14 June 2012

Printed in USA